



A comparative study on distressful events in affective disorder and normal control

Abstract

Background: Life events' stresses are concerned with situational encounters and the meaning that a person attaches to such encounters. It refers to our feeling; it is something of importance to us and is being jeopardised by events in our daily life, and the stressful life events are causally linked to a variety of undesirable effects which influence our performance and health. **Aim:** This study was planned for assessment and comparison of stressful life events between mood disorder and normal people. **Materials and methods:** In this study, total 90 participants (30 manic patients, 30 depressive patients, and 30 normal participants) were recruited and severity of symptoms was assessed by Young Mania Rating Scale and Beck Depression Inventory. Normal participants were screened by General Health Questionnaire. Presumptive Stressful Life Events Scale was used for both groups for assessment of stressful life events. **Findings and conclusion:** The present study results revealed that clinical group had higher score on stressful life events as compared to normal participants. Patients with depression had more stressful life events as compared to the mania and normal population. Overall, life events precede the mood symptoms' occurrence.

Keywords: Mania. Depression. Psychological Stress.

Sushma Rathee¹, Pradeep Kumar²

¹PhD Scholar, Department of Psychology, Maharshi Dayanand University, Rohtak, Haryana, India, ²Consultant Psychiatric Social Work, State Institute of Mental Health, University of Health Sciences, Rohtak, Haryana, India

Correspondence: Sushma Rathee, PhD Scholar, MPhil (Clinical Psychology), Department of Psychology, Maharshi Dayanand University, Rohtak-124001, Haryana, India. sushmaratheecp@gmail.com

Received: 1 June 2016

Revised: 20 August 2016

Accepted: 8 October 2016

Epub: 20 November 2016

DOI: 10.5958/2394-2061.2016.00044.6

Introduction

Depression is a mood disorder in which main symptoms are related with mood disturbance, thought process, sadness, loneliness, hopelessness, self-doubt, and guilt. Depression could be divided into two major categories in which first is known as mild depression. It is known as mild because it has effect on daily life but at minimal level. In major depression type of categories, symptoms influence person's ability to deal with daily life activities. Periods of depression usually last from a few weeks to many months, but a period lasting for several years is not uncommon.[1] In mania, most frequent symptoms include hyperactivity, reduced need for sleep, racing thoughts, irritability, elation, aggressiveness, and delusions of grandeur.

A person with pessimistic thinking is more vulnerable to depression and mood disorders when he or she experiences negative events in life.[1] Behavioural views of affective disorders focus on types of reinforcement by reducing negative consequences and increasing positive reinforcements. The emotional or psychodynamic perspective returns to early parent-child and early caregiver-child relationships. The emotional perspective again believes that disruptions and inadequacies in early child relationships open the door to mood and affective disorders. Sociocultural components include low self-esteem and insufficient support systems with relation to gender or socioeconomic factors.[2] Mathew *et al.*[3] conducted a study in which they included 46 manic patients and they assessed their life events during a period of six months before and after

illness started. In their study, they found stressful life events were significantly higher before the first episode compared to after episode, and they also found that life events before illness were significantly associated with male patient and with a younger age of onset.[3] A recent study done by Rodrigo *et al.* in Asian country indicated that symptoms of depression were significant in 37.5%; they also found that patients and their caregivers showed strain which were associated with interruption to work and disputes with relations.[4]

Methods

In the present study, sample comprised of 90 subjects (30 manic patients, 30 depressive patients, and 30 normal participants). Those patients fulfilling diagnosis criteria according to ICD-10 were included.[5] Their age range was between 20-50 years. Participants were able to read Hindi. Those having comorbid psychiatric illness or having bipolar disorder and presence of any major medical or neurological illness were excluded from the study.

Research design

The study was cross-sectional in nature and those participants fulfilling the criteria were selected.

Setting

The study was conducted in Psychiatry Department, Pt BD Sharma PGIMS, Rohtak.

Duration of the study

The study was conducted during May to October 2014.

Measures of the study

Socio-demographic proforma and clinical data

A proforma was specially designed for collecting information about socio-demographic and clinical variables of sample. It included information such as age, sex, residence, marital status, education, occupation, duration of current episode, history of present illness, number of previous depressive episodes, significant past psychiatric history, significant medical history, family history of medical illness, family history of psychiatric illness, significant personal history, and diagnosis.

Young Mania Rating Scale

Young Mania Rating Scale (YMRS) was developed by Young *et al.* in 1978 to assess the severity of manic symptoms.[6] YMRS consists of total 11 items. Beck Depression Inventory-II: Beck Depression Inventory-II (BDI-II) was developed by Beck *et al.* in 1996 to assess the level of depression.[7] This scale consists of 21 items. General Health Questionnaire (GHQ): General Health Questionnaire (GHQ) was used as a screening instrument to identify psychological distress among adults.[8] GHQ consists of 28 items. Presumptive Stressful Life Events Scale: Presumptive Stressful Life Events Scale (PSLES) was developed by Singh *et al.* in 1984.[9] This scale consists of 51 items which is divided into two categories, i.e. past and present.

Statistical analysis

Statistical Package for Social Sciences (SPSS-16) was used for analysis. Data was assessed using mean and standard deviation, independent samples *t*-test discrete variables for comparisons.

Results

The general characteristics of the participants such as sample size, age, sex, occupation, marital status, and education were equally distributed. Total 90 participants were selected (clinical population=60 [30 mania, 30 depression], normal population=30). In the descriptive analysis of mean and standard deviation of the socio-demographic details, both the clinical and normal groups were comparable.

Table 1 shows results of the stressful life events in the previous year, past life, and total score on PSLES between patients with mania, depression, and normal population. In the previous or one year before illness was started, in past years, and in total, stressful life events were found higher in clinical group as compared to normal group. Overall, total score showed that clinical population had significant higher stressful life events than normal participants.

In Table 2, *t*-test was calculated to compare the stressful life events in three conditions which was, in the recent one year, in past years of life, and total score on PSLES. Results showed significant difference in the total score that predicted

that life events which occurred during whole life influenced the psychological health.

Table 3 shows the results of various domains in PSLES between patients with mania, depression, and normal population. Clinical group have higher score in the domains of bereavement, education, family & social (manic), financial, health, legal (depressive), marital & sexual, and work of PSLES compared to normal control.

Discussion

In the present study, age range of the sample was 20-50 years. Most of the patients were diagnosed in early adulthood. Participants varied in education from illiterate to graduation in both groups; in clinical group, it was found that most of the patients either had no formal education or education up to secondary level. In sex ratio, affective disorder was found equally in both male and females. In this study, onset of illness was found in early 20's, and duration of illness was chronic and episodic in nature. Stressful life events in the previous one year, past, and total score was found significantly higher in the clinical group than normal group. Manic patients had higher score in past years on PSLES as compared to depressive patients and patients with depressive disorder had higher score in overall total score. Overall results showed that clinical population had significantly higher stressful life events than normal participants. In support of present study, literature has also found a positive association between life events and affective episodes.[10] A study conducted by Kumari and Jahan[11] showed that distressful life events were present in all three groups. However, life events of manic group were mainly related to social life, whereas life events of depressed and normal subjects were mostly related to personal life.[11] Hosang *et al.*[12] conducted a study in which they found stressful life events to be significantly associated with unipolar depression and bipolar disorder. In a recent study done by Koenders *et al.*,[13] they found that negative life events were significantly associated with both types of mood disorder.

Our study showed the results in the domains of PSLES which were family & social, financial, work, marital & sexual, health, education, legal, courtship & cohabitation, and bereavement. Descriptive analysis was done in which the results were found that clinical group had higher score in health, family & social, financial, work, marital & sexual, and bereavement in patients with mania whereas in depressive patients results were found that they had higher score on financial, work, marital & sexual, bereavement, legal, health, and education. Both clinical and normal

Table 1: Mean and standard deviation of stressful life events in previous year, past life, and total score between patients with mania, depression, and normal population

PSLES domain	Manic patients (N=30)	Depressive patients (N=30)	Normal control (N=30)
One year	238.83±73.57	264.10±79.72	211.63±106.91
Past years	176.40±50.49	148.60±67.12	128.67±73.91
Total	408.033±63.40	423.03±76.02	337.60±114.236

PSLES=Presumptive Stressful Life Events Scale

Table 2: t-test between patients with mania, depression and normal control

PSLES domain	Manic patients (t value) (N=30)	Depressive patients (t value) (N=30)	Degree of freedom	Significance (2-tailed)
One year	2.155	2.15	58	0.036
Past years	1.011	1.01	58	0.316
Total	2.77	2.88	58	0.005

PSLES=Presumptive Stressful Life Events Scale

Table 3: Descriptive statistics between patients with mania, depression, and normal control on various domains of PSLES

PSLES domain	Manic patients (N=30)	Depressive patients (N=30)	Normal control (N=30)
Bereavement	54.50±40.25	76.13±35.18	49.66±34.55
Courtship and Cohabitation	9.86±23.88	15.70±24.67	15.76±21.07
Education	23.56±29.77	25.83±27.22	23.73±22.55
Family and Social	129.40±77.67	97.73±57.61	110.50±71.86
Financial	23.30±39.23	42.77±35.31	12.77±23.55
Health	60.03±46.13	39.17±32.93	27.97±25.62
Legal	4.0±15.54	39.76±44.28	10.40±24.15
Marital and Sexual	17.40±49.72	19.27±43.15	7.13±18.90
Work	44.93±46.44	43.20±32.86	40.87±32.33

PSLES=Presumptive Stressful Life Events Scale

group had higher score in the family & social as compared to other domains; it means that family & social domain had significant role and influence in the life of both the groups. In favour of result of the present study, a study conducted by Proudfoot *et al.*[14] found that psychological stress acting as trigger was specifically associated with the onset of manic/hypomanic episodes/depressive episodes. To know about the role life events play in genesis of disorders, Medhi and Das[15] used the PSLES. Number of life events and total life events scores were significantly higher in depressive group than control group. Close association of bereavement and depression was observed.[15] Ghosh and Dutta[16] used PSLES to assess life event in major psychoses including mood disorder. Mood disorder patients involve both mania and major depression.[16] They found close relation of major life events with depression and mania, and specially mentioned bereavement as a life event.[16]

Limitation and future direction

Present study was conducted on a small sample size. Hence, result of the present study cannot be generalised. Structured diagnostic tool had not been used to confirm the diagnosis. Study was cross-sectional in nature. In future, this study can be done on a large sample. In the present study, we assumed that stressful life events were significant between patients of affective disorder and normal population. We found significant difference in the stressful life events between all three groups.

Conclusion

Overall significant difference was found between both groups in PSLES; clinical population had higher score than normal population and both groups had higher score in previous year of stressful life events. Life events appeared

before the occurrence of mood symptoms. In the study, significant difference was found in clinical as well as in between mania and depression also. Significant findings were found that both groups had higher score on the family & social domain of PSLES. Our study assessed these findings in Indian context.

References

1. Macneil CA, Hasty MK, Conus P, Berk M, Scott J. Bipolar disorder in young people: a psychological intervention manual. Cambridge: Cambridge University Press; 2010.
2. Estrin D, Sim I. Health care delivery. Open mHealth architecture: an engine for health care innovation. *Science*. 2010;330:759-60.
3. Mathew MR, Chandrasekaran R, Sivakumar V. A study of life events in mania. *J Affect Disord*. 1994;32:157-61.
4. Rodrigo C, Fernando T, Rajapakse S, De Silva V, Hanwella R. Caregiver strain and symptoms of depression among principal caregivers of patients with schizophrenia and bipolar affective disorder in Sri Lanka. *Int J Ment Health Syst*. 2013;7:2.
5. World Health Organization. The ICD-10 classification of mental and behavioural disorders: clinical descriptions and diagnostic guidelines. Geneva, World Health Organization; 1992.
6. Young RC, Biggs JT, Ziegler VE, Meyer DA. A rating scale for mania: reliability, validity and sensitivity. *Br J Psychiatry*. 1978;133:429-35.
7. Beck AT, Steer RA, Brown GK. BDI-II, Beck depression inventory: manual. San Antonio, Tex: Psychological Corp, Boston: Harcourt Brace; 1996.
8. Golderberg D, Williams P. A user's guide to the General Health Questionnaire. Windsor, UK: NFER-Nelson; 1988.
9. Singh G, Kaur D, Kaur H. Presumptive stressful life events scale (PSLES) - a new stressful life events scale for use in India. *Indian J Psychiatry*. 1984;26:107-14.
10. Alloy LB, Abramson LY, Urosevic S, Walshaw PD, Nusslock R, Neeren AM. The psychosocial context of bipolar disorder: environmental, cognitive, and developmental risk factors. *Clin Psychol Rev*. 2005;25:1043-75.
11. Kumari A, Jahan M. Distressful life events in affective disorder. *Journal of the Indian Academy of Applied Psychology*. 2006;32(3):193-200.

12. Hosang GM, Korszun A, Jones L, Jones I, McGuffin P, Farmer AE. Life-event specificity: bipolar disorder compared with unipolar depression. *Br J Psychiatry*. 2012;201:458-65.
13. Koenders MA, Giltay EJ, Spijker AT, Hoencamp E, Spinhoven P, Elzinga BM. Stressful life events in bipolar I and II disorder: cause or consequence of mood symptoms? *J Affect Disord*. 2014;161:55-64.
14. Proudfoot J, Whitton A, Parker G, Doran J, Manicavasagar V, Delmas K. Triggers of mania and depression in young adults with bipolar disorder. *J Affect Disord*. 2012;143:196-202.
15. Ghosh S, Dutta D. Study on role of life event in major psychoses. *Dysphrenia*. 2014;5:127-32.
16. Medhi D, Das PD. Clinical study on role of life events in genesis of neurotic disorders and depression. *Open J Psychiatry Allied Sci*. 2015;6:138-42.

Rathee S, Kumar P. A comparative study on distressful events in affective disorder and normal control. *Open J Psychiatry Allied Sci*. 2017;8:58-61. doi: 10.5958/2394-2061.2016.00044.6. Epub 2016 Nov 20.

Source of support: Nil. **Declaration of interest:** None.