



A case study of alcohol dependence syndrome with poor motivation and coping skills: the psychosocial perspective

Abstract

This case study is an attempt to assess the impact of psychiatric social work intervention in person with alcohol dependence. Psychiatric social work intervention (brief intervention) was provided to the client focusing on building motivation for change and strengthening commitment to change. It uses a single subject design and compares pre- and post-intervention baseline data with that following intervention. Semi-structured clinical and socio-demographic data sheet, family assessment proforma, and readiness to change questionnaires were administered to the client. The brief psychiatric social work intervention was provided to the client and family members. The attempt has been to bring out changes in motivation level and to enhance coping skill. After brief psychiatric social work intervention, knowledge regarding the illness was enhanced. The client motivation level was enhanced, family members have better understanding about client's illness, and interpersonal relationship has been improved.

Keywords: Alcohol Dependence. Coping Skills. Psychiatric Social Work Intervention.

U Harikrishnan¹, Arif Ali²

¹M. Phil, Department of Psychiatric Social Work, LGB Regional Institute of Mental Health, Tezpur, Assam, India, ²Assistant Professor, Department of Psychiatric Social Work, LGB Regional Institute of Mental Health, Tezpur Assam, India

Correspondence: U Harikrishnan, Department of Psychiatric Social Work, LGB Regional Institute of Mental Health, Tezpur, Assam, India.
hariarchal@gmail.com

Received: 27 April 2016

Revised: 23 October 2016

Accepted: 24 October 2016

Epub: 31 October 2016

DOI: 10.5958/2394-2061.2016.00037.9

Introduction

Alcoholism constitutes a vast syndrome of medical, economics, psychological, and social problems. Brief interventions for alcohol-related problems are time limited, structured, and directed toward a specific goal. Brief interventions can be used to change substance abuse behaviours and involve a variety of approaches, ranging from unstructured counselling and feedback to formal structured therapy.[1-6] Brief psychosocial interventions play an important role in determination of treatment outcomes in addiction [7-10]. It has been shown to improve client compliance to medication and the retention of clients in treatment. It has also been shown to increase alcohol abstinence rates and enhance quality of life in persons with alcohol dependence. Psychiatric social work plays a key role in working with those affected by alcohol addiction – from client's psychosocial assessment and intervention to working with the family of the addicted person. Social workers are part of the multidisciplinary team; they are directly involved in psychosocial intervention/brief intervention and recovery of addictions.

Methodology

It uses a single subject design and compares pre- and post-intervention baseline data with that following intervention. The attempt has been to bring out changes in motivation level.

Assessment

1. Semi-structured clinical and socio-demographic data sheet: A relevant sociodemographic and clinical detail was collected using this proforma.
2. Family assessment proforma: The interview was done with this client based on the family assessment proforma by Bhatti and Mathew.[11] The family assessment proforma is based on the family structure, leadership patterns, role and functioning, communication, cohesiveness, and adaptive patterns.
3. Readiness to Change Questionnaires (RCQ): The RCQ (treatment version)[12] is a 15-item self-report questionnaire that evaluates treatment-seeking individuals to three stages of change that is pre-contemplation, contemplation, and action stage of change. The RCQ (treatment version) is a 5-point Likert scale ranges from "strongly disagree" (-2) to "strongly agree" (+2). The internal consistency of scale range is between 0.60 (contemplation) and 0.70 (action). The scale's retest reliability ranges between 0.69 (contemplation) and 0.86 (action).

Case introduction

The index client was 42 years old, Hindu, married for 12 years from middle socioeconomic background, hailing from urban area of Kamrup district of Assam. The client was admitted in a hospital for alcohol-related problem.

Sources of information

The client himself, his mother, client's wife, and case record file were sources of information, which were reliable and adequate.

The reason for referral

The case was referred to the department of psychiatric social work for adequate psychosocial assessment and intervention.

Brief for clinical history

Index client Mr. BB was apparently well 15 years back. The client started drinking alcohol when he was 20 years old along with his friends at college. Initially, he was taking alcohol (30 ml) – two pegs, but then gradually his intake of alcohol had increased. In 2000, the pattern of alcohol intake was increased due to his job pressure. Mr. BB was involved in the private job. He started having decreased sleep and appetite, distress, irritability, and poor concentration. Due to the above reasons, he started drinking two to three large pegs (60 ml) daily at morning and evening, and he started giving excuses to his manager saying that he was not well. Since 2010 onward client's intake of alcohol had been increased (375 ml/day). Generally he used to take alcohol alone. Since the last three months, clients reported of decreased sleep, tremor, weakness, and restlessness. Due to the above reasons, the client came for treatment, diagnosed as F 10.2 (alcohol dependence syndrome) and was admitted in the hospital for further management.[13]

Family genogram

The client's family history (Figure 1): Client's father, 70 years old, worked as a government employee. Client's mother, she was 66 years old, homemaker. The client's second sibling was sister, 40 years old, married, and studied up to graduation. Third sibling was sister, 34 years old, married, and studied up to graduation. The relationship between client and family member was satisfactory earlier, currently they were having strained relationship due to client's alcoholic behaviour. The interaction patterns between client's parents were not adequate, client's father used to have argument and quarrel with his wife. The interaction between siblings was adequate toward client, they showed concerns and support toward




Figure 1: Family genogram

the client, and they only worried about the client's alcoholic behaviour.

Family dynamic

Some level of close and rigid boundaries was evident among the subsystems. The nominal and functional of the family was client's father. Decision was reached with mutual consultation and informed to the family members. In family, each member was performing their role adequately except the client due to his illness. In communication pattern, both verbal and non-verbal communication was present in the family. Positive reinforcement was present in the family and the support system was satisfactory in nature. We feeling was not present in the family. Client's siblings came to help his family whenever the need arose or at the time of crisis. Problem solving ability and coping strategy were found to be poor among the family.

Personal history

In personal history, the birth order of the client was the first among the three siblings. The client had a normal birth and development milestones were age appropriate and achieved normally. No major health problems were reported during the childhood. The client was an active child and there was no major behaviour problems in childhood. In educational history, the client started going to school at around four and half years of age. There were frequent changes in school, because of his father's nature of job. He was an average student, studied up to master in business administration. He had a healthy interaction with peer group. Frequent job changes were reported by the informants. The client married when he was 30 years old and spouse was 29 years at the time of marriage. She was staying with the client's family. In marital history, it was found that there was frequent quarrel, arguments, and conflict between the couple and marital satisfaction was not adequate because of client's alcoholic behaviour. Client's daughter was 11 years old, studying in sixth standard. The client was very much caring and loving toward his child.

Premorbid personality

Client was having well-adjusted premorbid personality.

Z-diagnosis

- Z 56 - Problems related to employment and unemployment
- Z 71 - Alcohol use
- Z 72.3 - Lack of physical exercise
- Z 72.4 - Inappropriate diet and eating habits
- Z 73.3 - Stress, not elsewhere classified.

Table 1: Pre and post-test in RCQ

RCQ	Pre-test	Post-test
Pre-contemplation stage	-8	-6
Contemplation	+7	+8
Action	+7	+8

RCQ: Readiness to Change Questionnaires

Table 1 shows the pre- and post-test of the clients; in pretest, it was found that motivation level was in precontemplation stage (score -8). After psychiatric social work intervention, post-test was done after one month of intervention and RCQ score suggested that motivation of client to quit alcohol had been enhanced. The client was found in the action phase (score +8).

Psychiatric social work intervention

For this case study, psychiatric social work intervention means brief intervention (psychoeducation, motivation enhancement therapy [MET], relapse prevention therapy, coping skills, supportive case work, admission counselling, and pre-discharge counselling) provided to the client and family members. The total number of sessions conducted was 14.

Process of intervention

Rapport establishment

Rapport establishment aims to maintain a good relationship with the client, and to assess the level of cooperation and participation of the client. During the discussion, the client was informed about the importance of therapy and benefit he would gain. Repeated reassurance and positive attitude toward the client made the session successful. The client was able to open up in the initial session itself; he expressed and shared about his drinking problem.

Family assessment and intervention

The family assessment was done with the client's wife to address the psychosocial issues of the family and also to investigate the level of knowledge regarding mental illness due to prolonged period of alcohol abuse by the client. On family assessment, it was found that due to client's alcoholic behaviour wife had reported depressive and anxiety symptoms, codependency was present, crying spells, and poor interpersonal relationship with client. In marital assessment, it was found that there was frequent quarrel and arguments between them. Supportive intervention was provided to the wife. Coping skill was taught to the client's wife. The therapist provided psychoeducation to the family members so that there was a better understanding toward the client's illness.

MET

MET evokes change in individuals. It is a systematic approach and is based on motivational psychology. MET is considered to produce change through mobilising the person's internal resources. Thus, the intervention is brief and used during the client's first contact although repeating it during subsequent sessions may prove helpful. It is particularly useful when contact with the client is limited to one or few sessions. The treatment outcome research supports MET as effective in producing change in problem drinkers.[14-18] The model for stages of behavioural change developed by Prochaska and DiClemente[19] can be utilised in understanding how people change their behaviour and in assessing the individual's readiness to change a particular behaviour. The model includes the following stages of change: precontemplation, contemplation, determination/preparation, action, and maintenance.

The index client was exposed to motivation enhancement and relapse prevention therapy. A baseline assessment on understanding the drinking pattern, abstinence period, locus of control, coping pattern, internal relationship problems, and client's attitude toward drinking was done. Throughout the MET sessions, the therapist recognised the client's individual efforts, appreciated his strengths, and feedbacks were provided. Further, in MET the client was provided warmth, advice, empowerment, and support.

Relapse prevention technique

Craving management skill

Craving management skills were taught to the client. "5 Ds", i.e. Delay, Deep breathing, Distraction, Drinking water, and Discussion was taught to the client to deal with the craving.

Coping skills

Coping skills' training was provided to the client with objectives to teach how to avoid high-risk situations, to recognise urge "triggers", and handling stressor.

Social group work interactive sessions

A group is a collection of individuals with similar problems. Group work intervention helps them to discuss and share their experiences with one another, and through this process, learns skills of coping, decision making, and problem solving. The goal of the group work was as follows:

- Accepting the fact that alcohol is a problem
- Recognising the existence of other problems related to alcohol dependency
- Becoming aware of and identifying feelings
- Recognising and managing relapse cues
- Identifying one's own cues and plan to handle
- Handling stressful situation
- Helping him changing his lifestyle
- Improving interpersonal relationship
- Learning new ways to respond to problems
- Assisting in maintaining abstinence.

The number of members in the group was eight with two psychiatric social work trainees. Each session for 45 minutes once in a week. The client's level of awareness was increased, and he got insight regarding his problem of drinking.

Predischarge counselling

A brief psychoeducation was given to the client regarding mental illness and negative effects of alcohol. The following areas were discussed:

- The nature of mental illness
- The need of regular medication
- The harmful effects of alcohol
- The importance of getting involved in productive work
- Dealing with stigma
- Follow-up.

Discharge counselling with client and family

Client and his wife attended the session. The session focused on the illness and treatment modalities, importance of regularity of medication, importance of regular follow-up, family involvement in treatment, importance of work engagement, adopting healthy lifestyle, and importance of family support.

Follow-up session

After the discharge, the client came for follow-up. Feedback was taken from the client and his wife. It was found that client was maintaining well and was following the instruction and advice provided to him in dealing with his alcoholic behaviour.

Discussion

In this case study it was found that after psychiatric social work intervention, there was difference in pre- and post-test scores in RCQ. In a study, Borah and Ali[20] based on psychiatric social work assessment and intervention to the person with alcohol dependence syndrome, focused on building motivation for change and strengthening commitment to change. They found that after the end of the therapy, the client developed better knowledge regarding the illness, motivation level was enhanced to action phase, and coping skills were enhanced to recover from relapse.[20] Brief psychosocial assessment and intervention have shown to be efficacious in treatment and also improving psychosocial functioning in individual with alcohol abuse and dependence.[21-30] A brief psychosocial intervention along with pharmacological treatment is necessary in the management of alcohol dependence to bring back to normal life. In the case study, it was found that client motivation level was enhanced and coping skills to deal with the problems were improved; further it was found that interpersonal relationship had been improved.

Conclusion

Psychiatric social work interventions for the treatment of alcohol and drug problems cover a diverse array of treatment interventions. These interventions generally focus on the individual (their beliefs, feelings, and behaviour), their social context, including family and community.

References

- Kristenson H, Ohlin H, Hultén-Nosslin MB, Trell E, Hood B. Identification and intervention of heavy drinking in middle-aged men: Results and follow-up of 24-60 months of long-term study with randomized controls. *Alcohol Clin Exp Res*. 1983;7:203-9.
- Chick J, Lloyd G, Crombie E. Counselling problem drinkers in medical wards: A controlled study. *Br Med J (Clin Res Ed)*. 1985;290:965-7.
- Persson J, Magnusson PH. Early intervention in clients with excessive consumption of alcohol: A controlled study. *Alcohol*. 1989;6:403-8.
- Zweben A, Fleming MF. Brief interventions for alcohol and drug problems. In: Tucker JA, Donovan DM, Marlatt GA, editors. *Changing Addictive Behaviour: bridging clinical and public health strategies*. New York: Guilford Press; 1999:251-82.
- Beich A, Gannik D, Malterud K. Screening and brief intervention for excessive alcohol use: Qualitative interview study of the experiences of general practitioners. *Br Med J*. 2002;325:870-5.
- Bray JW, Zarkin GA, Davis KL, Mitra D, Higgins-Biddle JC, Babor TF. The effect of screening and brief intervention for risky drinking on health care utilization in managed care organizations. *Med Care*. 2007;45:177-82.
- Center for Substance Abuse Treatment. *Brief Interventions and Brief Therapies for Substance Abuse Treatment Improvement Protocol 34*. HHS Publication No. [SMA] 99-3353. Rockville, MD: Substance Abuse and Mental Health Services Administration; 1999.
- Heather N. The public health and brief interventions for excessive alcohol consumption: The British experience. *Addict Behav*. 1996;21:857-68.
- Wutcke SE, Shiell A, Gomei MK, Conigrave KM. Cost effectiveness of brief interventions for reducing alcohol consumption. *Soc Sci Med*. 2001;52:863-70.
- Borsari B, Carey KB. Effects of a brief motivational intervention with college student drinkers. *J Consult Clin Psychol*. 2000;68:728-33.
- Bhatti RS, Mathew V. Family therapy in India. *Indian J Soc Psychiatry*. 1995;11:30-4.
- Heather N, Luce A, Peck D, Dunbar B, James I. The development of a treatment version of the readiness to change questionnaire. *Addict Res*. 1999;7:63-8.
- World Health Organization. *The ICD-10 classification of mental and behavioural disorders: clinical descriptions and diagnostic guidelines*. Geneva: World Health Organization; 1992.
- Miller WR. Motivational interviewing with problem drinkers. *Behav Psychother*. 1983;11:147-72.
- Miller WR. Motivation for treatment: a review with special emphasis on alcoholism. *Psychol Bull*. 1985;98:84-107.
- Miller WR. Rediscovering fire: small interventions, large effects. *Psychol Addict Behav*. 2000;14:6-18.
- Hettema J, Steele J, Miller WR. Motivational interviewing. *Annu Rev Clin Psychol*. 2005;1:91-111.
- Angami MA, Baruah A, Ahmed N. Motivational intervention among clients with alcohol dependence syndrome: a quasi experimental study in a selected de-addiction centre in Assam. *Dysphrenia*. 2014;5:91-7.
- Prochaska JO, DiClemente CC. Stages of change in the modification of problem behaviours. In: Hersen M, Eisler RM, Miller PM, editors. *Progress in Behaviour Modification*. Sycamore, IL: Sycamore Publishing Company; 1992:184-214.
- Borah S, Ali A. A case study of person with alcohol dependence syndrome with poor motivation. *Int Res J Soc Sci*. 2016;5:74-9.
- Elzerbi C, Donoghue K, Drummond C. A comparison of the efficacy of brief interventions to reduce hazardous and harmful alcohol consumption between European and non-European countries: A systematic review and meta-analysis of randomized controlled trials. *Addiction*. 2015;110:1081-90.
- Kaner EF, Beyer F, Dickinson HO, Pienaar E, Campbell F, Schlesinger C, *et al*. Effectiveness of brief alcohol interventions in primary care populations. *Cochrane Database Syst Rev*. 2007;CD004148.
- Witkiewitz K, Marlatt GA: *Therapist's Guide to Evidence-Based Relapse Prevention*. London: Academic Press; 2007.
- Marlatt GA, Witkiewitz K. Relapse prevention for alcohol and drug problems. In: Marlatt GA, Dennis MD, editors. *Relapse Prevention: Maintenance Strategies in the Treatment of Addictive Behaviours*. 2nd ed. New York: Guilford Press; 2005:1-44.
- Irvin JE, Bowers CA, Dunn ME, Wang MC. Efficacy of relapse prevention: A meta-analytic review. *J Consult Clin Psychol*. 1999;67:563-70.
- McCready BS. Alcohol use disorders and the Division 12 Task Force of the American Psychological Association. *Psychol Addict Behav*. 2000;14:267-76.
- Miller WR, Rollnick S. *Motivational Interviewing: Preparing People for Change*. 2nd ed. New York: Guilford Press; 2002.
- Huang YS, Tang TC, Lin CH, Yen CF. Effects of motivational enhancement therapy on readiness to change MDMA and methamphetamine use behaviours in Taiwanese adolescents. *Subst Use Misuse*. 2011;46:411-6.
- Dieperink E, Fuller B, Isenhardt C, McMaken K, Lenox R, Pocha C, *et al*. Efficacy of motivational enhancement therapy on alcohol use disorders in clients with chronic hepatitis C: A randomized controlled trial. *Addiction*. 2014;109:1869-77.
- Korte KJ, Schmidt NB. Motivational enhancement therapy reduces anxiety sensitivity. *Cogn Ther Res*. 2013;37:1140-50.
- Harikrishnan U, Ali A. A case study of alcohol dependence syndrome with poor motivation and coping skills: the psychosocial perspective. *Open J Psychiatry Allied Sci*. 2017;8:87-90. doi: 10.5958/2394-2061.2016.00037.9. Epub 2016 Oct 31.

Source of support: Nil. Declaration of interest: None.