



Dyadic adjustment and marital communication of persons with depression

Abstract

Background: One of the most important relationships between a man and a woman is marriage. Marriage is a social union as well as a legal contract between people that creates kinship. It is a primary institution of society. Marriage involves an emotional and legal commitment that is quite important in any adult life and even same in the persons with depression. Faulty marital relationship not only affects the couples but it inflicts longitudinal negative impact upon the entire family system and individual members of families. **Methodology:** The present study aims to compare the pattern of dyadic adjustment and marital communication between married individuals with depression and married healthy controls. Cross-sectional study design was used. Purposive sampling technique was used to collect the data for the study. The study group comprised of 30 married individuals with depression, whereas the other group consisted of 30 married healthy controls. Both the groups were examined on sociodemographic datasheet, the Dyadic Adjustment Scale (DAS), and the Marital Communication Inventory (MCI). **Result:** It was found that there were significant differences in the dyadic adjustment and marital communication of persons with depression compared to healthy controls, and there was no significant difference between males and females of depression patients in dyadic adjustment and marital communication. **Conclusion:** There is need to involve the spouse in the treatment process, and couple-based interventions will help for better marital adjustment and communication with depression patients compared to the healthy controls.

Keywords: Depressive Disorder. Marital Adjustment. Couple Communication.

**Prasad Kannekanti¹,
Dipanjan Bhattacharjee¹, Daya Ram²,
Hariom Pachori³**

¹Department of Psychiatric Social Work,
Central Institute of Psychiatry, Ranchi, India,
²Department of Psychiatry, Central Institute
of Psychiatry, Ranchi, India, ³Department
of Statistics, Central Institute of Psychiatry,
Ranchi, India

Correspondence: Dr. Prasad Kannekanti,
Assistant Professor, Department of Psychiatric
Social Work, Central Institute of Psychiatry,
Ranchi, India-834006.
reddysriprasad@gmail.com

Received: 30 April 2020

Revised: 8 February 2021

Accepted: 10 February 2021

Epub: 20 February 2021

DOI: 10.5958/2394-2061.2021.00025.2

INTRODUCTION

Depressive disorders are one of the major leading causes of disability due to their high prevalence rates. They will cost a huge burden financially as well due to lack of productivity.[1] Depression is one of the common mental disorders, and persons with depression have a higher risk of suicide rate than the general population. Risk factors of depression are manifold, but family and marital factors have been recognised as essential reasons. In 2010, depression was estimated as the second most common cause of years lived with disability (YLD) worldwide.[2]

Marriage is one of the basic social institutions in human society. It produces family and fulfils human society's needs in an organised way, which is the distinct feature from animal society. A healthy marital relationship must have following characteristics: a) positively oriented healthy communication between married couples, b) presence of reciprocal as well as complementary relationship, c) emotionality in relationship or presence of emotional attachment between couples, and d) mutual sense of responsibility.[3] The systemic transactional model (STM) explains that rather than the individualistic view of stress and coping, it is more of partners' mutual interdependent experiences. During interactions between

the couples, the dyad acts as the medium of sharing feelings, negotiating opinions, and coping with stressors.[4,5] The general consensus suggests that marriage has potential protective qualities against psychological problems including depression.[6,7] However, some past observations suggested that the psychological benefits of marriage largely depend on the interpersonal processes operating in the marital dyads.[8-10]

Globally, depressive disorders are one of the top three leading causes of YLD in 2017 for both genders.[11] Numerous studies had shown that psychiatric disorder has significant negative impacts on almost all aspects of marital life.[12-15] Not only marital life, perhaps, but the psychiatric disorder also has a negative impact on a person and his/her all spheres. Psychiatric disorders limit one's performance and abilities in every sphere of life. Some research indicates that marital problems and psychological disorders reciprocally influence one another.[16] Nevertheless, it is equally true that there is no simple explanation for the association between the poor quality of marriage and depression. Both poor marriage and depression form part of the complex system of interactions represented in the bi-directional relationship between marriage and depression.[17,18] In Indian society, marriage is considered one of the most important institutions

that has a significant role in the interplay of social norms and values. Marriage is a social bond and meets the intimate needs of human beings which guide the path for procreation and maintenance of the human race. Empirical evidence showed that men and women in satisfying marriages appear to be at lower risk for a psychiatric disorder than other segments of the population.[19-21] Several past empirical studies showed that in marital dyads with one partner with depression have some distinctive characteristics, such as paucity of verbal and nonverbal positivity, asymmetry in interaction patterns between couples, presence of a high degree of passivity/withdrawn relationships between partners, frequent occurrences of negative statements between the couples (e.g. complaints, negative self-statements), and negative reciprocity between them.[22,23] Couples with a depressed partner were found to be negative and less congenial in their marital interactions, and depressive symptoms also predict the weakening of the couple's marital adjustment and satisfaction.[22,24,25]

METHODS AND MATERIALS

The present study aimed to compare the pattern of dyadic adjustment and marital communication between married individuals with depression and married healthy controls. This was a cross-sectional study. Purposive sampling was used to collect the data from the outpatient department and psychosocial unit of the Central Institute of Psychiatry (CIP), Kanke, Ranchi, Jharkhand, India. This study was conducted with 60 samples, 30 married individuals with depression as study group and control group with 30 samples matched with the age, sex, duration of the marriage, and family income with the study group. Objectives of the study were to compare dyadic adjustment and marital communication among persons with depression and healthy controls, and to look at the difference between males and females suffering from depression. Inclusion criteria were persons with depression diagnosed according to the tenth revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-10),[26] married for at least two years living with the spouse in the age range of 20-50 years with a minimum of fifth class education. Healthy controls were taken after screening with the General Health Questionnaire (GHQ)-12;[27] persons who scored less than or equal to three were considered as healthy controls. Exclusion criteria were the presence of comorbid psychiatric, physical, or neurological problems. After obtaining permission from the institutional review board of CIP, consent was obtained from the respondents, data was collected, and all ethical

considerations were taken care of in the process of data collection.

Tools used in the study

Sociodemographic datasheet

Sociodemographic datasheet was prepared by the researcher to collect the basic sociodemographic details of the respondents.

General health questionnaire (GHQ)-12

GHQ-12[27] is a screening device for identifying minor mental health issues in the general population. It consists of 12 items and scored on a four-point Likert-type scale (from zero to three). The score was used to generate a total ranging from zero to 36. The positive items were corrected from zero (always) to three (never), and the negative ones from three (always) to zero (never). High scores indicate worse health. Internal consistency of this tool is excellent. A high degree of internal consistency is found for all the 12 items with Cronbach's alpha value of 0.37-0.79, while total score was 0.79.

Dyadic adjustment scale (DAS)

The Dyadic Adjustment Scale (DAS)[28] is a 32-item self-reported scale. It has been used in many studies to measure marital adjustment and satisfaction. It has four subscales: dyadic satisfaction, dyadic cohesion, dyadic consensus, and affectional expression. Scores range from zero to 151 on the total scale; a higher score indicates better adjustment. The tool consistently discriminates between distressed and non-distressed couples, and cut off scores of 97 and 100 have been used. Internal reliability estimates of 0.96 were found for DAS; Cronbach's alpha for this sample was 0.93.

Marital communication inventory (MCI)

The Marital Communication Inventory (MCI)[29] is a self-report measure of marital communication that assesses processes like spouse's ability to express themselves and their style of expressions. It is a 46-item inventory, arranged on a four-point rating scale with options of usually, sometimes, seldom, and never. The tool yields a single total score. Scores range from zero to 138. Higher scores indicate better communication. The split-half reliability was 0.93. The inventory could also reliably discriminate between individuals with good and poor marital communication. The tool has been widely used in marital research, evaluation of marital counselling, and family life education programmes. It has also been widely used to assess communication and therapy change in Indian couples.[30]

Table 1: Comparison of dyadic adjustment between study group and control group (N=60)

Domains of Dyadic Adjustment Scale (DAS)	Groups (N=60)		t (df=58)	p
	Study group	Control group		
	Mean±SD (n=30)	Mean±SD (n=30)		
Dyadic consensus	42.50±5.22	53.63±2.89	-10.210	<0.001
Dyadic satisfaction	31.26±4.83	38.56±2.73	-7.550	<0.001
Dyadic cohesion	14.06±2.19	18.50±2.56	-7.183	<0.001
Affectional expression	6.73±1.28	10.26±1.04	11.671	<0.001
Total DAS score	94.56±11.08	120.96±8.07	-10.546	<0.001

Statistical analysis

The statistical analysis was performed using Statistical Package for the Social Sciences (SPSS) 16.0. To compare the study group with healthy control individuals, chi-square and independent samples *t* test were used.

RESULTS

Table 1 shows the comparison of scores in different domains and aggregate of DAS between depressive patients and healthy controls. Very significant differences ($p < 0.001$) were observed between these two groups in both domain specific and aggregate scores of DAS.

Table 2 shows the marital communication between the study and control group. Very significant difference ($p < 0.001$) was observed between these two groups in the scores of MCI.

Table 3 shows the comparison of dyadic adjustment and marital communication of males and females of depression group. There was no difference found in the domains of dyadic consensus, dyadic satisfaction, dyadic cohesion, affectional expression, marital communication, comparison between males and females.

DISCUSSION

The aim of the present study was to compare the pattern of dyadic adjustment and marital communication between married individuals with depression and married healthy controls. After fulfilling the inclusion and exclusion criteria, the current study recruited 30 married persons with depression and 30 married healthy controls. The study group was recruited from outpatient department and psychosocial unit of CIP and healthy control group was taken from the hospital surroundings. The mean age of the study group of depressed patients is 35.33 ± 6.94 years and control group is 35.83 ± 4.69 years, consisted of 30 married healthy controls matched with the samples of the study group on sociodemographic parameters (e.g. age, sex, duration of the

marriage, and monthly income). In India, the proportion of disability was estimated to be 67% to 70% among major depressive disorder.[31] Disability can affect marital relationships; the current dyadic adjustment scores and marital communication were found to be below cut-off scores among persons with depression.

Comparison of the dyadic adjustment of the depressed patients and the healthy controls on various domains of DAS,[28] i.e. dyadic consensus, dyadic satisfaction, dyadic cohesion, and affectional expression was made. Very significant differences ($p < 0.001$) were seen between patients' group and healthy control group in all domains of DAS as well as the total score of DAS. This finding is consistent with numerous studies done in the past in relation to dyadic communication and adjustment of depressed patients.[14,32,33] In the present study, individuals with depression showed problems (low scores in all areas of DAS in comparison to healthy controls) in dyadic adjustment because of their illness, which might cause marital problems in them. Empirical evidence had shown in the past that marital problems and depression are known to be among the most frequent problems for which adults seek treatment in a mental health facility.[34,35] Being depressed is often proved to be an antagonistic factor for marital life and it may cause significant distress to spouses of the depressed people.[36,37] This way present study has been in consonance with previous studies mentioned earlier.

In Asian countries, one of the most prominent aspects of couple's relationship is marital communication.[38,39] In the present study, marital communication has also been found to be problematic in depressed patients as depressed patients had scored significantly lower than healthy controls on MCI. Past studies showed that poor marital quality (characterised by shallow and insipid communications between spouses, lower presence of emotionality in communicational styles and contents, and overall faultiness in spousal communication) might lead to depression and depression may lead to poor marital quality, and it is probable that, when both are present, they reinforce each other. So, this way present study is also in consonance with previous studies that communication pattern as well as resources in married depressed people, is problematic because of inadequacy and inaccuracy.[37,40] That is why in the present study, married depressed people scored significantly lower than healthy controls. The quality of interpersonal relationship between the couple is also important while handling married depressive patients; these issues should not be neglected for holistic care. Longitudinal studies have shown that couple problems and depressive

Table 2: Comparison of marital communication between study group and normal group (N=60)

Score of Marital Communication Inventory	Groups (N=60)		t (df=58)	p
	Study group	Control group		
	Mean±SD (n=30)	Mean±SD (n=30)		
	83.23±9.34	119.86±4.38	-19.436	<0.001

Table 3: Comparison of dyadic adjustment and marital communication of males and females of depression group (N=30)

Variables	Males Mean±SD n=15	Females Mean±SD n=15	t (df=28)	p
Dyadic consensus	43.20±4.70	41.80±5.77	0.728	0.473
Dyadic satisfaction	32.20±2.45	30.33±5.88	1.133	0.267
Dyadic cohesion	14.26±1.96	13.86±2.47	0.492	0.626
Affectional expression	6.93±1.22	6.53±1.35	0.849	0.403
Marital communication	83.26±7.98	83.20±10.82	0.019	0.985

symptoms influence each other; either depressive symptoms predict marital issues or marital issues can lead to depression or vice versa.[41] Comparison of dyadic adjustment and marital communication of males and females of depression group (N=30) has shown that there was no difference found in the domains of dyadic consensus, dyadic satisfaction, dyadic cohesion, affectional expression, and marital communication between males and females. It reveals that depression affects both marital communication and adjustment for both genders.

Limitations

1. The present study was conducted with small sample size and purposive sampling design left scope for potential selection bias in the study.
2. The healthy controls' sociodemographic parameters varied from the study group as the patient group approaching for treatment had a diverse demographic structure while healthy controls were the subjects of the hospital surroundings only.

Conclusion

In the current study, the dyadic adjustment and marital communication were found to be low as reported by persons with depression compared to healthy control, which indicates the need for couple-based psychosocial interventions for the married persons with depressive disorders. If depressive symptoms act as risk factors in marital relationship, couple-based interventions will address depressive symptoms as well couple relationship issues. Couple-based indigenous psychosocial interventions are needed with this population.

ACKNOWLEDGEMENTS

We thank all the staff of Central Institute of Psychiatry (CIP), Kanke, Ranchi, Jharkhand, India for the support and cooperation during this study.

AUTHOR CONTRIBUTIONS

PK, DR, and DB designed the study; PK collected data independently and wrote the manuscript with the help of DB and DR; HP performed all the statistical tests; DB and DR supervised the findings of the work; DR provided critical feedback and helped shape the research work; all authors discussed the results and contributed to the final manuscript.

REFERENCES

1. Donohue JM, Pincus HA. Reducing the societal burden of depression: a review of economic costs, quality of care and effects of treatment. *Pharmacoeconomics*. 2007;25:7-24.
2. Ferrari AJ, Charlson FJ, Norman RE, Patten SB, Freedman G, Murray CJ, *et al*. Burden of depressive disorders by country, sex, age, and year: findings from the global burden of disease study 2010. *PLoS Med*. 2013;10:e1001547.
3. Barker P, Chang J. Basic family therapy. Chichester, West Sussex, UK: John Wiley & Sons; 2013.
4. Bodenmann G, Randall AK, Falconier MK. Coping in couples: the systemic transactional model (STM). In: Falconier MK, Randall AK, Bodenmann G, editors. *Couples coping with stress: a cross-cultural perspective*. New York, NY: Routledge; 2016:5-22.
5. Liu YC, Fang SY, Lin SH, Lin MF. Couple-based interventions for improving dyadic adjustment in people with cancer and their spousal partners. *Cochrane Database Syst Rev*. 2019;2019:CD013324.
6. Frech A, Williams K. Depression and the psychological benefits of entering marriage. *J Health Soc Behav*. 2007;48:149-63.
7. Waite LJ, Gallagher M. The case for marriage: why married people are happier, healthier, and better off financially. New York: Random House Digital; 2001.
8. Thomeer MB, Umberson D, Pudrovskaya T. Marital processes around depression: a gendered and relational perspective. *Soc Ment Health*. 2013;3:151-69.
9. Hawkins DN, Booth A. Unhappily ever after: effects of long-term, low-quality marriages on well-being. *Soc Forces*. 2005;84:451-71.
10. Williams K. Has the future of marriage arrived? A contemporary examination of gender, marriage, and psychological well-being. *J Health Soc Behav*. 2003;44:470-87.
11. GBD 2017 Disease and Injury Incidence and Prevalence Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 354 diseases and injuries for 195 countries and territories, 1990-2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet*. 2018;392:1789-858. Erratum in: *Lancet*. 2019;393:e44.
12. Murray CJL, Lopez AD, World Health Organization, World Bank & Harvard School of Public Health. The Global burden of disease: a comprehensive assessment of mortality and disability from diseases, injuries, and risk factors in 1990 and projected to 2020: summary [Internet]. World Health Organization; 1996 [cited 2020 Apr 30]. Available from: <https://apps.who.int/iris/handle/10665/41864>
13. Horwitz AV, White HR. The relationship of cohabitation and mental health: a study of a young adult cohort. *J Marriage Fam*. 1998;60:505-14.
14. Coyne JC, Benazon NR. Not agent blue: effects of marital functioning on depression and implications for treatment. In: Beach SRH, editor. *Marital and family processes in depression: a scientific foundation for clinical practice*. Washington, DC: American Psychological Association; 2001:25-43.
15. Isaac R, Shah A. Sex roles and marital adjustment in Indian couples. *Int J Soc Psychiatry*. 2004;50:129-41.
16. Gilligan M, Sutor JJ, Pillemer K. Estrangement between mothers and adult children: the role of norms and values. *J Marriage Fam*. 2015;77:908-20.
17. Coyne JC. Toward an interactional description of depression. *Psychiatry*. 1976;39:28-40.
18. Birtchnell J. Depression and family relationships. A study of young, married women on a London housing estate. *Br J Psychiatry*. 1988;153:758-69.
19. Birtchnell J. Negative modes of relating, marital quality and depression. *Br J Psychiatry*. 1991;158:648-57.
20. Weiss RL, Aved BM. Marital satisfaction and depression as predictors of physical health status. *J Consult Clin Psychol*. 1978;46:1379-84.
21. Weissman MM, Gammon GD, John K, Merikangas KR, Warner V, Prusoff BA, *et al*. Children of depressed parents. Increased psychopathology and early onset of major depression. *Arch Gen Psychiatry*. 1987;44:847-53.
22. Johnson SL, Jacob T. Marital interactions of depressed men and women. *J Consult Clin Psychol*. 1997;65:15-23.
23. Gabriel B, Beach SR, Bodenmann G. Depression, marital satisfaction and communication in couples: investigating gender differences. *Behav Ther*. 2010;41:306-16.
24. McCabe SB, Gotlib IH. Interactions of couples with and without a depressed spouse: self-report and observations of problem-solving situations. *J Soc Pers Relat*. 1993;10:589-99.
25. Halford WK, Doss BD. New frontiers in the treatment of couples. *Int J Cogn Ther*. 2016;9:124-39.
26. World Health Organization. The ICD-10 classification of mental and behavioural disorders: clinical descriptions and diagnostic guidelines. Geneva: World Health Organization; 1992.
27. Goldberg D, Williams P. A user's guide to the General Health.

- Questionnaire. Windsor, UK: NFER-Nelson; 1988.
28. Spanier GB. Measuring dyadic adjustment: new scales for assessing the quality of marriage and similar dyads. *J Marriage Fam.* 1976;38:15-28.
 29. Bienvenu MJ. Measurement of marital communication. *Fam Coord.* 1970;19:26-31.
 30. Mayamma MC, Sathyavathi K. Disturbances in communication and marital disharmony in neurotics. *Indian J Psychiatry.* 1985;27:315-9.
 31. Gururaj G, Varghese M, Benegal V, Rao GN, Pathak K, Singh LK, *et al.* National mental health survey of India, 2015-16: summary. Bengaluru: National Institute of Mental Health and Neurosciences; 2016.
 32. Umberson D, Williams K, Powers DA, Liu H, Needham B. You make me sick: marital quality and health over the life course. *J Health Soc Behav.* 2006;47:1-16.
 33. Proulx CM, Helms HM, Buehler C. Marital quality and personal well-being: a meta-analysis. *J Marriage Fam.* 2007;69:576-93.
 34. Beach SR, O'Leary KD. Treating depression in the context of marital discord: outcome and predictors of response for marital therapy vs. cognitive therapy. *Behav Ther.* 1992;23:507-28.
 35. Beach SR, O'Leary KD. Dysphoria and marital discord: are dysphoric individuals at risk for marital maladjustment? *J Marital Fam Ther.* 1993;19:355-68.
 36. Beach SR, Fincham FD, Katz J. Marital therapy in the treatment of depression: toward a third generation of therapy and research. *Clin Psychol Rev.* 1998;18:635-61.
 37. Benazon NR, Coyne JC. Living with a depressed spouse. *J Fam Psychol.* 2000;14:71-9.
 38. Karimi R, Bakhtiyari M, Masjedi Arani A. Protective factors of marital stability in long-term marriage globally: a systematic review. *Epidemiol Health.* 2019;41:e2019023.
 39. Sorokowski P, Randall AK, Groyecka A, Frackowiak T, Cantarero K, Hilpert P, *et al.* Marital satisfaction, sex, age, marriage duration, religion, number of children, economic status, education, and collectivistic values: data from 33 countries. *Front Psychol.* 2017;8:1199. Erratum in: *Front Psychol.* 2017;8:1728.
 40. Whisman MA, Uebelacker LA, Tolejko N, Chatav Y, McKelvie M. Marital discord and well-being in older adults: is the association confounded by personality? *Psychol Aging.* 2006;21:626-31.
 41. Kronmüller KT, Backenstrass M, Victor D, Postelnicu I, Schenkenbach C, Joest K, *et al.* Quality of marital relationship and depression: results of a 10-year prospective follow-up study. *J Affect Disord.* 2011;128:64-71.

Kannekanti P, Bhattacharjee D, Ram D, Pachori H. Dyadic adjustment and marital communication of persons with depression. *Open J Psychiatry Allied Sci.* 2021;12:115-9. doi: 10.5958/2394-2061.2021.00025.2. Epub 2021 Feb 20.

Source of support: Nil. **Declaration of interest:** None.