



Open Journal of Psychiatry & Allied Sciences (OJPAS®), January-June 2022, Volume 13 Issue 1

Page no.	Error	Correction
Title: OJPAS® this issue		
2	1. Nukala S, Vinnakota A, Garapati VA, Singiseti S, Chilukuri S, Sanapala V, Nambaru LR . COVID-19-related stigma towards doctors in India: an online cross-sectional study. <i>Open J Psychiatry Allied Sci.</i> 2022;13:21-5. doi: 10.5958/2394-2061.2022.00005.2. Epub 2022 Jan 29.	1. Nukala S, Vinnakota A, Garapati VA, Singiseti S, Chilukuri S, Sanapala V, et al . COVID-19-related stigma towards doctors in India: an online cross-sectional study. <i>Open J Psychiatry Allied Sci.</i> 2022;13:21-5. doi: 10.5958/2394-2061.2022.00005.2. Epub 2022 Jan 29.
Title: Conceptual framework of psychiatric care in India: moving from community psychiatry to public psychiatry		
3	The purpose of lunatic asylums was to segregate persons with mental illness (PWMI) from the community and not for the purpose of their treatment.	The purpose of lunatic asylums was to segregate PWMI from the community and not for the purpose of their treatment.
4	Psychiatric camp (extension/ satellite/ outreach clinic)[10] Conducted on the pre-decided date and place. Moved from informal non-healthcare settings (temple, school, gram panchayat, etc.)[9] to formal healthcare settings (nearby primary health centres, taluk hospitals, private clinics, etc.)[10,11]	Psychiatric camp (extension/ satellite/ outreach clinic)[9] Conducted on the pre-decided date and place. Moved from informal non-healthcare settings (temple, school, gram panchayat, etc.)[10] to formal healthcare settings (nearby primary health centres, taluk hospitals, private clinics, etc.)[9,11]
5	Even though in DMHP , there is mention of awareness creation, stigma reduction, early detection, and rehabilitation of PWMI, these goals took a back seat, and the focus remained on the treatment of severe mental illnesses.	Even though in District Mental Health Programme (DMHP) , there is mention of awareness creation, stigma reduction, early detection, and rehabilitation of PWMI, these goals took a back seat, and the focus remained on the treatment of severe mental illnesses.
6	Further, the 'clinical psychiatry' can be conveniently classified four-tier based hierarchy of healthcare delivery system as 'primary care psychiatry' at primary health centres by primary care doctors (MBBS), secondary care/ general hospital psychiatry at district/ taluk hospitals/ multi-specialty hospitals/ private clinics by psychiatry specialists, tertiary care/ general hospital psychiatry at medical colleges, and quaternary care psychiatry at apex medical/ mental health institutes.	Further, the 'clinical psychiatry' can be conveniently classified four-tier based hierarchy of healthcare delivery system as 'primary care psychiatry' at primary health centres by primary care doctors (Bachelor of Medicine, Bachelor of Surgery [MBBS]), secondary care/ general hospital psychiatry at district/ taluk hospitals/ multi-specialty hospitals/ private clinics by psychiatry specialists, tertiary care/ general hospital psychiatry at medical colleges, and quaternary care psychiatry at apex medical/ mental health institutes.
8	9. Sidana A. Community psychiatry in India: where we stand? J Mental Health Hum Behav. 2018;23:4-11. 10. Burns T. Community psychiatry at the National Institute of Mental Health and Neuro Sciences, India: easier than working in Manchester. Psychiatr Bull. 1998;22:180-2.	9. Burns T. Community psychiatry at the National Institute of Mental Health and Neuro Sciences, India: easier than working in Manchester. Psychiatr Bull. 1998;22:180-2. 10. Sidana A. Community psychiatry in India: where we stand? J Mental Health Hum Behav. 2018;23:4-11.
Title: Family burden among caregivers of alcohol dependence: a cross-sectional study		
14	Leff J, Wig NN, Bedi H, Menon DK, Kuipers L, Korten A, et al. Relatives' expressed emotion and the course of schizophrenia in Chandigarh. A two-year follow-up of a first-contact sample. <i>Br J Psychiatry.</i> 1990;156:351-6.	Leff J, Wig NN, Bedi H, Menon DK, Kuipers L, Korten A, et al. Relatives' expressed emotion and the course of schizophrenia in Chandigarh. A two-year follow-up of a first-contact sample. <i>Br J Psychiatry.</i> 1990;156:351-6.
Title: Evaluation of attitude and regard towards mental illness in comparison with chronic medical conditions among final year nursing students		
16	Among the study group, 11.46% of the students had experienced mental illness at some point in time and sought professional help and 22.29% had family history of mental illness.	Among the study group, 11.46% of the students had experienced mental illness at some point in time and sought professional help , and 22.29% had family history of mental illness.
Title: COVID-19-related stigma towards doctors in India: an online cross-sectional study		
23	This probably is reflective of the fact that professional satisfaction of the doctor goes much beyond perceived stigma, and the appreciation and gratitude expressed by patients predominates , and on the whole, providing them with a rewarding experience.	This probably is reflective of the fact that professional satisfaction of the doctor goes much beyond perceived stigma, and the appreciation and gratitude expressed by patients predominate , and on the whole, providing them with a rewarding experience.

Open Journal of Psychiatry & Allied Sciences (OJPAS®), January-June 2022, Volume 13 Issue 1

Page no.	Error	Correction
24	Stigma and discrimination towards frontline workers may be mitigated by raising awareness on print and digital media and highlighting the role of doctors and other healthcare workers in containing the pandemic.	Stigma and discrimination towards frontline workers may be mitigated by raising awareness on print and digital media , and highlighting the role of doctors and other healthcare workers in containing the pandemic.
25	Nukala S, Vinnakota A, Garapati VA, Singiseti S, Chilukuri S, Sanapala V, Nambaru LR . COVID-19-related stigma towards doctors in India: an online cross-sectional study. Open J Psychiatry Allied Sci. 2022;13:21-5. doi: 10.5958/2394-2061.2022.00005.2. Epub 2022 Jan 29.	Nukala S, Vinnakota A, Garapati VA, Singiseti S, Chilukuri S, Sanapala V, et al . COVID-19-related stigma towards doctors in India: an online cross-sectional study. Open J Psychiatry Allied Sci. 2022;13:21-5. doi: 10.5958/2394-2061.2022.00005.2. Epub 2022 Jan 29.
Title: Entomophagy and coprophagy in alcohol use disorder: a rare psychopathology		
27	Non-contrast computed tomography (NCCT) head and x-ray chest was normal.	Non-contrast computed tomography (NCCT) head and x-ray chest were normal.
Title: Efficacy of brief cognitive behavioural intervention for the treatment of psychosis: a case report		
28	From the clinical interview, it was found that the patient was suffering from auditory hallucinations, delusion of control, and thought broadcasting among the positive symptoms of schizophrenia and had blunt affect and social withdrawal among the negative symptoms.	From the clinical interview, it was found that the patient was suffering from auditory hallucinations, delusion of control, and thought broadcasting among the positive symptoms of schizophrenia , and had blunt affect and social withdrawal among the negative symptoms.
30	8. National Collaborating Centre for Mental Health (UK). Psychosis and schizophrenia in adults: treatment and management: Updated edition 2014 [Internet]. London: National Institute for Health and Care Excellence (UK); 2014 [cited 2020 Apr 15]. Available from: https://www.nice.org.uk/guidance/cg178/evidence/full-guideline-490503565	8. National Collaborating Centre for Mental Health (UK). Psychosis and schizophrenia in adults: treatment and management: updated edition 2014 [Internet]. London: National Institute for Health and Care Excellence (UK); 2014 [cited 2020 Apr 15]. Available from: https://www.nice.org.uk/guidance/cg178/evidence/full-guideline-490503565
	10. Halder S, Mahato A. Cognitive remediation therapy in chronic schizophrenia. Chronic Mental Illness and the Changing Scope of Intervention Strategies, Diagnosis, and Treatment . IGI Global; 2017.	10. Halder S, Mahato A. Cognitive remediation therapy in chronic schizophrenia. Chronic mental illness and the changing scope of intervention strategies, diagnosis, and treatment . IGI Global; 2017.
Title: Our experience with implementation of manual augmentative and alternative communication (AAC) in adult with intellectual disability: a story of success		
31	The client expressed needs through vocalisation and bilabial sounds and his pragmatic skills included greetings, social smile.	The client expressed needs through vocalisation and bilabial sounds , and his pragmatic skills included greetings, social smile.

- Editor-in-Chief,

Open Journal of Psychiatry & Allied Sciences (OJPAS®)